

Accepted for publication in the Astrophysical Journal Letters

An empirical initial-final mass relation from hot, massive white dwarfs in NGC 2168 (M35)

Kurtis A. Williams

Steward Observatory

933 N. Cherry Ave., Tucson, AZ 85721

`kurtis@as.arizona.edu`

M. Bolte

UCO/Lick Observatory

University of California, Santa Cruz, CA 95064

`bolte@ucolick.org`

and

Detlev Koester

Institut für Theoretische Physik und Astrophysik

Universität Kiel, 24098 Kiel, Germany

`koester@astrophysik.uni-kiel.de`

ABSTRACT

The relation between the zero-age main sequence mass of a star and its white-dwarf remnant (the initial-final mass relation) is a powerful tool for exploration of mass loss processes during stellar evolution. We present an empirical derivation of the initial-final mass relation based on spectroscopic analysis of seven massive white dwarfs in NGC 2168 (M35). Using an internally consistent data set, we show that the resultant white dwarf mass increases monotonically with progenitor mass for masses greater than $4M_{\odot}$, one of the first open clusters to show this trend. We also find two massive white dwarfs foreground to the cluster that are otherwise consistent with cluster membership. These white dwarfs can be

explained as former cluster members moving steadily away from the cluster at speeds of $\lesssim 0.5$ km/s since their formation and may provide the first direct evidence of the loss of white dwarfs from open clusters. Based on these data alone, we constrain the upper mass limit of WD progenitors to be $\geq 5.8M_{\odot}$ at the 90% confidence level for a cluster age of 150 Myr.

Subject headings: white dwarfs — open clusters and associations: individual (NGC 2168)

1. Introduction

White dwarfs (WDs) are the final state of stellar evolution for the vast majority of intermediate- and low-mass stars. The upper mass limit of WD progenitor stars, M_{crit} , is also the lower mass limit of core-collapse supernova progenitors. The current best observational estimate for M_{crit} comes from spectroscopic analysis of WDs in the open cluster NGC 2516 (Koester & Reimers 1996) and is $M_{\text{crit}} \approx 8M_{\odot} \pm 2M_{\odot}$. Due to the steepness of the initial-mass function (IMF), however, this range results in a factor of ≈ 2 uncertainty in the number of supernovae and duration of supernova-driven winds resulting from bursts of star formation. This uncertainty in turn has a large impact on understanding the star-formation rate in galaxies (e.g. Somerville & Primack 1999), the evolution of starbursts (e.g. Leitherer et al. 1999), and the fate of low-mass dwarf galaxies at early times (e.g. Dekel & Silk 1986).

The best observational constraints on M_{crit} are obtained from studies of WD populations in open clusters with ages $\lesssim 150$ Myr (Williams 2002). NGC 2168 (M35) is one of the richest, compact and nearby open clusters in this age range, with age determinations ranging from ~ 100 Myr (von Hippel et al. 2000) to ~ 200 Myr (Sung & Bessell 1999). The WD cooling sequence of NGC 2168 has been discussed often in the literature, with recent photometric analyses by von Hippel et al. (2002) and Kalirai et al. (2003).

With spectroscopy of cluster WD candidates, it is possible to determine unambiguously if the objects are WDs and, for the bona fide WDs, to determine T_{eff} and $\log g$, and to derive the cooling age (τ_{cool}), mass and luminosity. For those WDs with cooling ages smaller than the cluster age and distance modulus consistent with cluster membership, subtraction of the WD cooling age from the age of the open cluster results in the lifetime of the progenitor star, and stellar evolutionary models can then be used to determine the progenitor star mass. Reimers & Koester (1988) applied this algorithm to two NGC 2168 WDs candidates identified on photographic plates by Romanishin & Angel (1980); however, their signal-to-noise was too low to determine the WD properties precisely.

As part of our ongoing program to identify and spectroscopically analyze the WD populations of open clusters, the Lick-Arizona White Dwarf Survey (LAWDS, Williams et al. 2004), we have obtained high signal-to-noise spectra of eight candidate massive WDs in NGC 2168. Six or seven of the observed WDs are hot, high-mass WDs likely to be cluster members. This sample doubles the number of known open cluster WDs with high-mass progenitors. A detailed photometric and spectroscopic analysis of the entire WD population of this cluster will be presented in a later paper. In this Letter we present the constraints on M_{crit} and the upper end of the initial-final mass relation for WDs based on the high-mass WDs already analyzed.

2. Observations & Analysis

UBV imaging of NGC 2168 was obtained in September 2002 with the Lick Observatory Shane 3-m reflector and the PFCam prime-focus imager; additional *UBV* imaging of a larger field center on the cluster was obtained in January 2004 with the KPNO 4m MOSAIC camera. PSF-fitting photometry was obtained using the DAOPHOT II program (Stetson 1987). Candidate WDs were selected by their blue excess in *UBV* color space. Figure 1 shows the color-magnitude diagram of all objects detected in *U*, *B*, and *V* across the entire MOSAIC field. Several very blue, faint objects are observed in the diagram; these are our candidate WDs. All four WD candidates of Romanishin & Angel (1980) are recovered, as is the WD candidate in von Hippel et al. (2002). Astrometry and photometry for the WD candidates in this Letter are presented in Table 1.

Spectroscopic observations of selected WD candidates were obtained with the blue camera of the LRIS spectrograph on the Keck I 10-m telescope (Oke et al. 1995). A 1''-wide longslit at parallactic angle was used with the 400 l mm⁻¹, 3400Å blaze grism for a resulting spectroscopic resolution of $\sim 6\text{\AA}$. The spectra were extracted and a relative spectrophotometric calibration applied using standard *IRAF*¹ routines.

The T_{eff} and $\log g$ were determined for each WD using simultaneous Balmer-line fitting (Bergeron, Saffer, & Liebert 1992). The model spectra are updated versions of those in Finley, Koester, & Basri (1997). The WD evolutionary models of Wood (1995) were used to calculate the mass (M_{WD}) and cooling age (τ_{WD}) of each WD. A distance modulus to each WD was measured by comparing the observed *V*-magnitude to the absolute magnitude

¹*IRAF* is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation

M_V calculated from the best-fitting model atmosphere and the appropriate WD cooling model. Errors in the fits were determined empirically by adding the noise measured for each spectrum to the best-fitting model spectrum convolved with the instrumental response. These simulated spectra were fit by the same method; nine iterations were used to calculate the scatter in T_{eff} , $\log g$, M_{WD} and τ_{WD} . The fitting procedure is discussed in depth in an upcoming paper on the open clusters NGC 6633 and NGC 7063 (Williams et al. 2004). The atmospheric fits and derived WD masses and ages are given in Table 2. The Balmer line fits are shown in Figure 2.

A systematic error in the fits of the hottest ($T_{\text{eff}} \gtrsim 50000\text{K}$) WDs became apparent, as it was not possible to simultaneously fit all the Balmer lines. In these cases, the fits were limited to $\text{H}\beta$, $\text{H}\gamma$, and $\text{H}\delta$. For LAWDS 22, no satisfactory convergence was achieved with these limited fits; the best-fit models are used in the analysis. WDs with $T_{\text{eff}} \gtrsim 50000\text{K}$ are known to exhibit metals in the atmosphere and NLTE effects (e.g. Napiwotzki 1992; Holberg, Barstow, & Sion 1998), neither of which are included in the models.

The progenitor mass for each WD was calculated by subtracting the WD cooling age from the cluster age. The age difference is the total lifetime of the progenitor star. The lifetimes of stars as a function of mass and metallicity are calculated from the stellar isochrones of Girardi et al. (2002). The progenitor mass for each WD likely to be a cluster member is given in Table 3 for an assumed cluster age of 150 Myr and for three different stellar evolutionary models: $Z = 0.008$ and $Z = 0.019$, both with modest convective overshoot, and $Z = 0.019$ without convective overshoot. Upper and lower errors are for 1σ differences in τ_{WD} .

3. Discussion

All eight of the observed WDs are much more massive than the typical WD mass of $\approx 0.56M_{\odot}$ (Bergeron, Saffer, & Liebert 1992), and five of the WDs have apparent distance moduli $(m - M)_V \approx 10.5$. It is therefore reasonable to assume that at least five of these objects are members of NGC 2168. LAWDS 11 is almost certainly *not* a cluster member. Its age is likely older than that of the cluster as a whole, and while the uncertainties in the spectral fits leave open the possibility that it is younger (which would require it to be hotter), the observed colors are more consistent with the cooler (and older) interpretation. The distance modulus of this WD is foreground to the cluster by little more than 1σ , but the cooler temperature favored by the $B - V$ color again favors the foreground interpretation.

LAWDS 15 has a mass and age consistent with cluster membership, but the calculated

distance modulus is inconsistent with that of the other WDs by $\sim 4\sigma_{M_V}$. Assuming that the distance modulus is correct and that there is no difference in reddening between LAWDS 15 and NGC 2168, LAWDS 15 is ~ 185 pc closer than the cluster. Based on the spectral fits, LAWDS 15 has a cooling age of ~ 50 Myr. If the WD has been moving away from the cluster since its formation at a steady rate of only 0.4 km/s, it will have covered this distance. Therefore, it is possible that LAWDS 15 was once a cluster member and has escaped the cluster.

The likelihood that a massive, hot WD would be found foreground along the line of sight to the cluster can be estimated from the luminosity function in Figure 16 of Liebert, Bergeron, & Holberg (2004). The luminosity function gives the space density of WDs with $M > 0.8M_\odot$ and $\tau_{\text{cool}} \leq 100$ Myr as $\sim 10^{-5.3} \text{ pc}^{-3} 0.5 \text{ mag}^{-1}$. This results in an estimated 0.1 hot, massive WDs in the $\sim 30' \times 30'$ MOSAIC field to a distance of 1 kpc. Therefore it is unlikely, but not impossible, that LAWDS 15 is a field WD. Based on these arguments, we will consider LAWDS 15 to be a cluster WD for this discussion. For similar arguments, we retain LAWDS 6 ($\sim 2\sigma_{M_V}$ closer than the other cluster WDs) as a likely cluster member.

Figure 3 shows the initial-final mass relation of these seven cluster members, along with that of WDs from Hyades, Praesepe and the Pleiades (Claver et al. 2001) and from NGC 2516 (Koester & Reimers 1996). Also shown are theoretical and semi-empirical data from plots in Claver et al. (2001) and sources therein. For the sake of consistency, the initial and final masses of each WD from the literature have been re-determined using our WD models and the published T_{eff} and $\log g$.

From the figure, it can be seen that the NGC 2168 WDs form a monotonic sequence of more massive WDs originating from more massive progenitors. This conclusion is robust, as changes in the assumed age of NGC 2168 do not affect the relative positions of the points, only the absolute initial masses. Figure 3 also shows that the NGC 2168 initial-final mass relation agrees with the empirical relation derived from previous clusters. This need not be expected *a priori*, as NGC 2168 has a significantly lower metallicity than the other clusters in the diagram. Model core masses decrease with increasing metallicity, and the efficiency of mass loss processes could change with metallicity.

LAWDS 15 and LAWDS 6, the potential escaped cluster WDs discussed above, fit the observed initial-final mass relation, providing additional evidence that they are cluster members. If these WDs are indeed escaped cluster members, they are crucial pieces of evidence that WDs can receive velocity kicks during mass loss and perhaps explain the observed deficit of WDs in other open clusters (see Williams 2004, and references therein). This and other potential explanations for these objects (e.g. binarity) will be discussed more fully in the later paper on WDs in NGC 2168.

The WD in Fig. 3 with the apparently low final mass is LAWDS 22. As mentioned above and visible in Fig. 2, the spectral fitting did not converge satisfactorily, despite the high signal-to-noise of the spectrum. This star is a close visual double with a redder companion ($V = 19.12$, $B - V = 1.27$, $U - B = 0.93$) $2''$ to the north. While resolved, this double is close enough that the spectrum of LAWDS 22 is likely contaminated by light from the neighbor star, resulting in the unsatisfactory fit. Light from the neighboring star may also be contaminating the photometric colors of LAWDS 22, which would also explain why the star lies redward of the $1M_{\odot}$ cooling track in Fig. 1. Given its measured T_{eff} , LAWDS 22 likely suffers from an extension of the systematic issue in the high- T_{eff} WD spectral fits described above. Other explanations for the location of this point could include magnetic fields (although no splitting is observed) or a low-mass, unresolved companion, but contamination by the neighboring star seems to be the most likely cause.

Based on the NGC 2168 data alone, it is possible to place lower limits on the value of M_{crit} . Making the assumption that errors in the WD ages in Table 2 are Gaussian, we calculate that the oldest cluster WD ages are $\log \tau_{\text{cool}} \geq 7.76$ with 90% confidence. For $\log \tau_{\text{cl}} = 8.15$ and $Z = 0.008$, this corresponds to $M_{\text{crit}} \geq 5.81M_{\odot}$ at a 90% confidence level. This value is in agreement with that obtained by Koester & Reimers (1996).

Improved constraints on the initial-final mass relation, including its intrinsic scatter and metallicity dependence, require improvements on existing observations. First and foremost, the ages of open clusters such as NGC 2168 must be determined to higher precision. Alternatives to main-sequence fitting such as lithium depletion studies or activity/rotation studies may provide the necessary constraints. Second, large samples of WDs from individual open clusters are needed to reduce the effect of systematics (such as errors in assumed ages) that plague the comparison of inter-cluster samples. The WD sample presented here is a start to that end, and the initial-final mass relation derived from these stars alone provides dramatic confirmation of the existence of an initial-final mass relation, an idea that was strongly hinted at from previous open cluster studies and from other theoretical and observational work. Assuming that the majority of the remaining massive NGC 2168 WD candidates are cluster WDs, planned spectroscopic observations of these objects will soon result in a sample of nearly a dozen of WDs originating from a single stellar population, permitting, for the first time, studies of the intrinsic dispersion of the initial-final mass relation at high masses.

M.B. and K.W. are grateful for support for this project in the form of the National Science Foundation AST-0307492. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

The authors would like to thank Matt Wood for freely providing the WD evolutionary models and for many useful comments regarding this paper. The authors also thank the anonymous referee for a well-considered report that resulted in several improvements to this Letter.

Some of the data presented herein were obtained at the W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

REFERENCES

- Bergeron, P., Saffer, R. A., & Liebert, J. 1992, *ApJ*, 394, 228
- Claver, C. F., Liebert, J., Bergeron, P., & Koester, D. 2001, *ApJ*, 563, 987
- Dekel, A. & Silk, J. 1986, *ApJ*, 303, 39
- Finley, D. S., Koester, D., & Basri, G. 1997, *ApJ*, 488, 375
- Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, *A&AS*, 141, 371
- Girardi, L., Bertelli, G., Bressan, A., Chiosi, C., Groenewegen, M. A. T., Marigo, P., Salasnich, B., & Weiss, A. 2002, *A&A*, 391, 195
- Holberg, J. B., Barstow, M. A., & Sion, E. M. 1998, *ApJS*, 119, 207
- Kalirai, J. S., Fahlman, G. G., Richer, H. B., & Ventura, P. 2003, *AJ*, 126, 1402
- Koester, D. & Reimers, D. 1996, *A&A*, 313, 810
- Leitherer, C., et al. 1999, *ApJS*, 123, 3
- Liebert, J., Bergeron, P., & Holberg, J. B. 2004, *ApJS*, in press (astro-ph/0406657)
- McCook, G. P. & Sion, E. M. 1999, *ApJS*, 121, 1
- Napiwotzki, R. 1992, *Lecture Notes in Physics*, Berlin Springer Verlag, 401, 310

- Oke, J. B., et al. 1995, *PASP*, 107, 375
- Reimers, D. & Koester, D. 1988, *A&A*, 202, 77
- Romanishin, W. & Angel, J. R. P. 1980, *ApJ*, 235, 992
- Somerville, R. S. & Primack, J. R. 1999, *MNRAS*, 310, 1087
- Stetson, P. B. 1987, *PASP*, 99, 191
- Sung, H. & Bessell, M. S. 1999, *MNRAS*, 306, 361
- von Hippel, T., Kozhurina-Platais, V., Platais, I., Demarque, P., & Sarajedini, A. 2000, *ASP Conf. Ser. 198: Stellar Clusters and Associations: Convection, Rotation, and Dynamos*, 75
- von Hippel, T., Steinhauer, A., Sarajedini, A., & Deliyannis, C. P. 2002, *AJ*, 124, 1555
- Weidemann, V. 2000, *A&A*, 363, 647
- Williams, K. A. 2002, Ph.D. Thesis
- Williams, K. A. 2004, *ApJ*, 601, 1067
- Williams, K. A., Bolte, M., Koester, D., & Wood, M. A. 2004, in preparation
- Wood, M. A. 1995, *LNP Vol. 443: White Dwarfs*, 41

Table 1. WDs in the field of NGC 2168, photometry.

LAWDS ID	McCook & Sion Designation ^a	RA (J2000)	Dec (J2000)	V	σ_V	$B-V$	σ_{B-V}	Previous References
NGC 2168:LAWDS 1	WD J0608+242	6:08:38.79	24:15:06.9	20.989	0.019	-0.035	0.028	1
NGC 2168:LAWDS 2	WD J0608+241	6:08:42.30	24:10:17.7	21.569	0.032	0.061	0.044	
NGC 2168:LAWDS 5	WD J0609+244.1	6:09:11.54	24:27:20.9	20.065	0.017	-0.128	0.024	1,2
NGC 2168:LAWDS 6	WD J0609+244.2	6:09:23.48	24:27:22.0	19.863	0.016	-0.128	0.023	1,2
NGC 2168:LAWDS 11	WD J0609+241	6:09:42.79	24:11:05.4	21.198	0.025	0.110	0.037	
NGC 2168:LAWDS 15	WD J0609+240	6:09:11.63	24:02:38.5	20.785	0.022	-0.039	0.032	
NGC 2168:LAWDS 22	WD J0608+245	6:08:24.65	24:33:47.6	19.657	0.016	0.008	0.023	
NGC 2168:LAWDS 27	WD J0609+243	6:09:06.26	24:19:25.3	21.398	0.026	0.090	0.039	3

^aFormat as in McCook & Sion (1999)

References. — (1) Romanishin & Angel (1980), (2) Reimers & Koester (1988), (3) von Hippel et al. (2002)

Table 2. WDs in the field of NGC 2168, spectral fits.

ID	T_{eff}	dT_{eff}	$\log g$	$d \log g$	M_{WD}	dM_{WD}	$\log(\tau_{\text{WD}})$	$d \log(\tau_{\text{WD}})$	M_V	dM_V	$(m - M)_V$
LAWDS 1	32400	512	8.40	0.125	0.888	0.075	7.386	0.208	10.454	0.215	10.535
LAWDS 2	31700	1800	8.74	0.191	1.072	0.102	7.929	0.291	11.132	0.328	10.437
LAWDS 5	52600	1160	8.24	0.095	0.824	0.051	6.158	0.025	9.542	0.184	10.523
LAWDS 6	55200	897	8.28	0.065	0.851	0.036	6.094	0.027	9.564	0.108	10.299
LAWDS 11	19900	2792	8.48	0.367	0.921	0.213	8.288	0.463	11.534	0.769	9.664
LAWDS 15	29900	318	8.48	0.060	0.934	0.037	7.693	0.089	10.754	0.111	10.031
LAWDS 22	54400	1203	8.04	0.121	0.721	0.060	6.169	0.054	9.155	0.219	10.502
LAWDS 27	30900	500	8.58	0.164	0.995	0.086	7.760	0.220	10.866	0.310	10.532

Table 3. Progenitor Masses for NGC 2168 WDs.

ID	$Z = 0.008$ ($M_{\odot}$)	$Z = 0.019$ ($M_{\odot}$)	$Z = 0.019^a$ ($M_{\odot}$)
LAWDS 1	$5.00^{+0.31}_{-0.16}$	$5.08^{+0.29}_{-0.15}$	$4.86^{+0.28}_{-0.14}$
LAWDS 2	$7.02^{+\infty}_{-1.60}$	$6.96^{+\infty}_{-1.49}$	$6.63^{+\infty}_{-1.40}$
LAWDS 5	$4.628^{+0.001}_{-0.001}$	$4.734^{+0.002}_{-0.001}$	$4.540^{+0.001}_{-0.001}$
LAWDS 6	$4.625^{+0.001}_{-0.001}$	$4.732^{+0.001}_{-0.001}$	$4.537^{+0.001}_{-0.010}$
LAWDS 15	$5.57^{+0.34}_{-0.24}$	$5.62^{+0.31}_{-0.22}$	$5.37^{+0.30}_{-0.21}$
LAWDS 22	$4.629^{+0.003}_{-0.003}$	$4.735^{+0.003}_{-0.002}$	$4.540^{+0.003}_{-0.002}$
LAWDS 27	$5.81^{+1.99}_{-0.61}$	$5.84^{+1.81}_{-0.56}$	$5.59^{+1.46}_{-0.54}$

^aNo convective overshoot




Fig. 1.— $B-V$, V color-magnitude diagram for NGC 2168. Squares indicate WDs presented in this study, both cluster members (filled) and the non-member (open). Solid curves indicate cooling curves for WDs with masses of $0.4M_{\odot}$ (top curve), $0.7M_{\odot}$, and $1.0M_{\odot}$ (bottom curve) at the distance and reddening of NGC 2168. The dotted line indicates the location of WDs with $\log \tau_{cool} = 8.15$, the assumed age of NGC 2168.




Fig. 2.— Balmer line fits for WD candidates in NGC 2168. The figures show the Balmer lines from H β (bottom) to H9 (top), with the curves showing the best-fit models.




Fig. 3.— Initial-final mass relation. Points with error bars are WDs from this study; other points are from the Hyades and Praesepe (Claver et al. 2001, open triangles), NGC 2516 (Koester 1996, open squares) and the Pleiad WD (Claver et al. 2001, upper limit). The starred point with error bars is LAWDS 22, discussed in the text. Curves represent the theoretical initial-final mass relation from Girardi et al. (2000) (solid), the core mass at the first thermal pulse from the same models (dashed), and the quasi-empirical relation from Weidemann (2000) (dotted line). Slight horizontal offsets have been applied to the three WDs with $M_{\text{init}} \approx 4.6 M_{\odot}$ for the sake of clarity. Statistical error bars in the initial masses for these three WDs are smaller than the points.